



## EXPERTISE-BASED EXPERTS IMPORTANCE WEIGHTS IN ADVERSE JUDGMENT

Evy Herowati<sup>1,2</sup>, Udisubakti Ciptomulyono<sup>1</sup>, Joniarto Parung<sup>2</sup> and Suparno<sup>1</sup>

<sup>1</sup>Industrial Engineering, Institute of Technology Sepuluh Nopember, Surabaya, Indonesia

<sup>2</sup>Industrial Engineering, University of Surabaya, Surabaya, Indonesia

E-Mail: [evy@ubaya.ac.id](mailto:evy@ubaya.ac.id)

### ABSTRACT

The objective of this research was to propose the use of expertise levels of experts to determine the experts' importance weights since there has been no research that determines the 'importance weight' using the expertise level as a whole. The significance of this research was the integration of three concepts, namely: the *expert's expertise level*, FPR's *Additive Consistency* and the *Induced-OWA operator* to obtain the expert's importance weight in adverse judgment situation. The Expertise level of an expert in adverse judgment situation is determined by his/her own assessment on a set of alternatives and defined as 'the ability to differentiate consistently' and expressed as the ratio between Discrimination and Inconsistency. The experts provided their preferences using FPR (Fuzzy Preference Relations) since FPR has Additive Consistency property to replicate each element of FPR matrix. Experts were sorted according to their expertise level and the experts' importance weights followed the OWA (Ordered Weighted Averaging) operator's weights which were determined by parameterization using Basic Unit-Interval Increasing Monotonic functions. The experts' importance weights model illustrated by a numerical example, and it concluded that the higher the expert's expertise level, the higher his/her importance weight.

**Keywords:** additive consistency, expertise, fuzzy preference relations, importance weight, induced OWA operator.

### INTRODUCTION

The expertise level of decision maker greatly affects the quality of the resulting decision [1]. The decision quality made by the experts presumed better than decision quality made by the non-experts as an expert have the ability to think differently [1-3]. This is because the inherent ability of the experts enables them to understand problems in more detail and depth so that the experts can distinguish the various aspects of the situation that is usually overlooked by the non-expert [4].

Related to decision that requires the Decision Maker (DM) expertise in Group Decision Making (Decision Making with more than one DM), the DM individual assessment needs to be explored and group decision is taken based on the integration of individual assessments into the group assessment, by performing aggregation of DM assessments mathematically [5]. One important factor that must be considered in the aggregation process is the importance weight of each DM. The magnitude of the DM's importance weight influence decisions. If a DM assessing an alternative with a high score and this DM gets a high importance weight, then this alternative would get a high total score and most likely has a high opportunity to be selected as the best alternative. Therefore, to improve the decision quality, the DMs' importance weights should be determined based on their expertise level.

Some researchers defined the expertise level as "the ability to differentiate consistently" through the evaluation of his/her assessment level on alternatives [6-7]. The level assessment on alternatives is called the *adverse judgment* [8]. Shanteau (2002) stated the experts as those who can differentiate between similar but not exactly the same, cases and repeat their assessments

consistently. They formulated the expertise level as the ratio between Discrimination ability and Inconsistency [6-7]. The drawback of this formulation found in the Inconsistency measurement [9]. Measuring Inconsistency required repetition; consequently the experts need to assess the same cases more than once. Assessing the same cases more than once is very difficult to do independently without being influenced by previous assessments. This is the reason of the need for adjustment to the formula implementation in determining the expertise level.

A number of researches have been proposed in determining the DMs importance weights and can be categorized into 2 groups: direct evaluation to DMs and evaluation to the DMs' assessment level. The DMs' importance weights determination through direct evaluation to DMs consists of a 'supra DM' who assessed the DM then gave weight to each DM [10-12] and 'a group of DM' who assessed each member in the DM group [13-15]. The direct evaluation methods could potentially lead to decision bias due to the assessors' subjectivity, the assessors' difficulty in assessing other DM and popularity effect (a person who has been recognized by peers usually assumed more expert; but it is possible that the 'assumed less expert' would be the creator of new knowledge [6]). While the determination of the DMs' importance weight through evaluation of their assessment on a set of alternatives can be classified into the determination of importance weights based on maximum consensus that could be achieved in the group [16-18], minimum deviation of DM individual opinion to the group opinion [19-21], minimum distance from the DM individual opinion to group opinion [22-27] and consistency of DM assessment on alternatives [8, 28-30]. In general, the methods of determining the DM importance weights in the



adverse judgment situation are more objective than direct evaluation to the DMs. These methods had certain property: the closer the DM's individual opinion on group average opinion, the greater his/her importance weight regardless of his/her expertise level. In particular, there were methods of importance weight determination based on the DM's consistency evaluation in pairwise comparison FPR (Fuzzy Preference Relation). These methods assumed that the more consistent the DM preferences, the more relevant the DM opinion and resulting the higher the DM importance weight. These consistency-based methods are more objective compared with the other methods, but these methods have not considered the expertise level as a whole.

In this paper, the DM importance weight is determined based on the expertise level as a whole as the ability to differentiate consistently and expressed by the ratio between Discrimination and Inconsistency and the difficulty in repetition is replaced by estimation using Additive Consistency (AC) of FPR. The DM provides preference in the form of pairwise comparisons FPR where FPR has the Additive Consistency (AC) properties that used to estimate repetition. The experts then sorted by their expertise level and the experts' importance weights are associated with the OWA (Ordered Weighted Averaging) operator's weights which are determined by parameterization using Basic Unit-Interval Increasing Monotonic functions (BUM).

To do this, this paper is organised as follows. Following the first Section, it is discussed the concepts used to obtain the DM's Importance weight, namely the expertise level of expert, the AC property of FPR and the Induced OWA Operator. Next, the methodology to obtain the *expertise-based experts importance weights* is discussed and followed by the applicability test of the methodology using a numerical example. Finally, in the last Section we drew our conclusions.

## LITERATURE REVIEWS

There are three important concepts used in this research to obtain the DM's Importance weight, namely the experts' expertise level, the AC property of FPR and the Induced OWA Operator.

### The Expertise Level of Experts

An expert usually has some backgrounds in certain fields and recognized by his/her peers [31]. Shanteau *et al.* (2002) determine the expertise level of an expert based on his/her assessment level (adverse judgment). They argued that only people who can differentiate between similar but not exactly the same, cases and repeat their judgment consistently, considered as an expert [6]. Therefore there are two requirements necessary for determining the expertise level, namely the Discrimination ability and the Inconsistency and expressed in CWS - Index as shown in eqn (1), (2) and (3):

$$\text{CWS-Index} = \frac{\text{Discrimination}}{\text{Inconsistency}} = \frac{\text{standard deviation of different alternatives' values}}{\text{standard deviation of the same alternative's values}} \quad (1)$$

$$\text{Discrimination} = \sqrt{\frac{\sum_{j=1}^n r(M_j - GM)^2}{n-1}} \quad (2)$$

$$\text{Inconsistency} = \sqrt{\frac{\sum_{j=1}^n \sum_{i=1}^r (M_{ij} - M_j)^2}{n(r-1)}} \quad (3)$$

Where

$r$	:	The number of replications
$M_j$	:	The average of individual values for case-j
$GM$	:	Grand mean of all individual values
$n$	:	The number of different cases
$M_{ij}$	:	The individual value for replication-i case-j

In order to measure the expertise level of the experts, the evaluated experts were asked to elicitate their evaluation more than once. Only those who have a high level of Discrimination ability and low level of Inconsistency can be classified as expert and obtain a high value of CWS-Index. Unfortunately, this method required repetition and this repetition are very hard to do independently without being affected by previous evaluation.

### The Additive Consistency of FPR

The Decision Makers could use a variety of evaluation formats, among others, is FPR. FPR is one of the most widely used evaluation format to provide evaluation in Group Decision Making (GDM) [16, 32-33] since FPR can be used as tools on aggregating individual opinions into a group opinion [34].

Suppose that a group of Decision Makers  $E = \{e_1, e_2, \dots, e_m\}$ ,  $m \geq 2$  evaluate a finite set of alternatives  $X = \{x_1, x_2, \dots, x_n\}$ ,  $n \geq 2$  by using pairwise comparisons FPR  $P \subset X \times X$  having a membership function  $\mu_p : X \times X \rightarrow [0,1]$  and represented by means of the  $n \times n$  matrix  $P = (p_{ij})$  [35].  $p_{ij}$  is the preference degree of alternative  $x_i$  over  $x_j$ .  $p_{ij} = 1/2$  means indifference between  $x_i$  and  $x_j$ ,  $p_{ij} > 1/2$  means  $x_i$  is preferred to  $x_j$ .

AC property of FPR among three alternatives  $x_i$ ,  $x_j$  and  $x_k$  [36] are as follows:

$$(p_{ij} - 0.5) + (p_{jk} - 0.5) = (p_{ik} - 0.5) \quad \forall i, j, k = 1, 2, \dots, n \quad (4)$$

$$p_{ij} + p_{jk} + p_{ki} = \frac{3}{2} \quad \forall i, j, k = 1, 2, \dots, n \quad (5)$$



The AC property yields a relationship between the preferences and we can obtain estimated values by using 3 different formulas as follows [28]:

$$ep_{ik}^{j1} = p_{ij} + p_{jk} - \frac{1}{2}, j \neq i, k \quad (6)$$

$$ep_{ik}^{j2} = p_{jk} - p_{ji} + \frac{1}{2}, j \neq i, k \quad (7)$$

$$ep_{ik}^{j3} = p_{ij} - p_{kj} + \frac{1}{2}, j \neq i, k \quad (8)$$

in which:

$ep_{ik}^{j1}$  : Estimation of  $p_{ik}$  using the first formula

$ep_{ik}^{j2}$  : Estimation of  $p_{ik}$  using the second formula

$ep_{ik}^{j3}$  : Estimation of  $p_{ik}$  using the third formula

For every element of the matrix FPR  $p_{ij}$ , these formulas produce as many as  $3 \times (n-2)$  replications (since there are 3 formulas and  $j \neq i, k$ ). These replications allow AC property be used to complete the incomplete FPR [28-29, 37-41] and measure someone's consistency level in providing assessment [28-29, 34, 39-40, 42] in which the consistency level then be used to determine the importance weight of each expert. The drawbacks are previous studies only considered the Consistency level, but not the Discrimination, thus have not covered the whole concept of expertise level as proposed by Shanteau *et al.*, (2002).

### The Induced OWA Operator

The OWA operator is an aggregation operator proposed by Yager (1988) in which the order of the arguments have primary role in the aggregation process [43]. An  $n$ -dimensional OWA Operator is a mapping  $F: I^n \rightarrow I$  defined as  $F_w(a_1, a_2, \dots, a_n) = \sum_{j=1}^n w_j b_j$  where  $b_j$  is the  $j$ th largest element in the set of input arguments  $(a_1, a_2, \dots, a_n)$  and  $w_j$  is the order weights satisfy  $w_j \geq 0$  and  $\sum_{j=1}^n w_j = 1$  [43]. In this case the input arguments are ordered according to their own values.

In the Induced OWA Operator, the ordering of the input arguments are based upon the order inducing variable [44]. An  $n$  dimensional Induced OWA Operator Is a mapping  $F: I^n \rightarrow I$  defined as  $F_w(\langle u_1, a_1 \rangle, \langle u_2, a_2 \rangle, \dots, \langle u_n, a_n \rangle) = \sum_{j=1}^n w_j b_j$  where  $u_i$  is called the order inducing variable and  $a_i$  is called input argument,  $w_j$  is the order weights and  $w_j \geq 0$  and  $\sum_{j=1}^n w_j = 1$  and  $b_j$  is the input argument value of the pair having the  $j$ -th largest value for the order inducing variable. An important issue in using the OWA operator or the Induced OWA operators in aggregation process is the

issue of obtaining the OWA weights. Yager (1996) proposed that the OWA weights can be parameterized by BUM  $Q: [0,1] \rightarrow [0,1]$  having the properties:

$Q(0) = 0; Q(1) = 1; Q(x) \geq Q(y)$  if  $x \geq y$  and the OWA weights  $w_j$  are as follows [45-46]:

$$w_j = Q(R_j) - Q(R_{j-1}) \quad (9)$$

where  $R_j > R_{j-1}$ , and the obtained weights satisfy  $w_j \geq 0$

and  $\sum_{j=1}^n w_j = 1$ . The BUM is associated with the

accumulation of DM importance weight. Since BUM is an increasing Monotonic Function as illustrated in Figure-1, then the individual DM importance weight can't be negative or  $w_j \geq 0$ .

Another important property of this function is the maximum value of BUM is one, so the accumulation of the total DM importance weights also 1. Yager (1988) proposed a particular form of BUM Function as  $Q(R) = R^\alpha$ ,  $\alpha$  positive parameter [43].




Figure-1. The OWA Weights from Basic Unit Interval Increasing Monotonic Function.

### THE PROPOSED METHOD

This research uses expertise level as 'the ability to differentiate consistently' and expressed as ratio between Discrimination and Inconsistency and the experts provide evaluation using pairwise comparison FPR. AC property of FPR enables us to get the replications without asking the DMs to repeat their evaluation and the result of this step is CWS-Index. Based on this Index, we obtain the rank of the DMs based on their assessment level [9, 47]. The next step is obtaining the DMs importance weights based on the Induced OWA weights and BUM function as illustrated in Figure-2.

The expertise-level expressed by the CWS-Index in logarithmic function and used as the order inducing variable. The expertise-level are used as the  $R$ -variable in determined the importance weights.

### NUMERICAL EXAMPLE


In order to show the applicability of the proposed methods, we provide a numerical example to illustrate



*Expertise-based experts' importance weight.* Suppose there are 5 experts expressed in  $E = \{e_1, e_2, e_3, e_4, e_5\}$  and

asked to provide assessment on a set of 4 alternatives  $X = \{x_1, x_2, x_3, x_4\}$ .

The data of experts' assessment are as follows [47]:



**Figure-2.** The Proposed Method.

$$R_1 = \begin{bmatrix} 0.5 & 0.4 & 0.4 & 0.4 \\ 0.6 & 0.5 & 0.4 & 0.7 \\ 0.6 & 0.6 & 0.5 & 0.7 \\ 0.6 & 0.3 & 0.3 & 0.5 \end{bmatrix}, \quad R_2 = \begin{bmatrix} 0.5 & 0.3 & 0.4 & 0.45 \\ 0.7 & 0.5 & 0.65 & 0.6 \\ 0.6 & 0.35 & 0.5 & 0.45 \\ 0.55 & 0.4 & 0.55 & 0.5 \end{bmatrix},$$

$$R_3 = \begin{bmatrix} 0.5 & 0.9 & 0.6 & 0.7 \\ 0.1 & 0.5 & 0.3 & 0.4 \\ 0.4 & 0.7 & 0.5 & 0.6 \\ 0.3 & 0.6 & 0.4 & 0.5 \end{bmatrix}, \quad R_4 = \begin{bmatrix} 0.5 & 0.43 & 0.64 & 0.71 \\ 0.57 & 0.5 & 0.43 & 0.36 \\ 0.36 & 0.57 & 0.5 & 0.21 \\ 0.29 & 0.64 & 0.79 & 0.5 \end{bmatrix},$$

$$R_5 = \begin{bmatrix} 0.5 & 0.7 & 0.8 & 0.6 \\ 0.3 & 0.5 & 0.65 & 0.55 \\ 0.2 & 0.35 & 0.5 & 0.4 \\ 0.4 & 0.45 & 0.6 & 0.5 \end{bmatrix}$$

The estimation of each matrix element by using formula 1, 2 and 3 will generate 6 estimated values. For example the estimated value of all matrix  $P_4$  elements is presented in Table-1. For each element of the matrix  $P_4$  there are 7 values ( $r = 7$ ), i.e. 6 estimated values and 1 real

value and we can calculate Discrimination and Inconsistency value.

$$\text{Discrimination} = \sqrt{\frac{\sum_{j=1}^n r(M_j - GM)^2}{n-1}} = \sqrt{\frac{1.54057}{(12-1)}} = 0.3742.$$

$$\text{Inconsistency} = \sqrt{\frac{\sum_{j=1}^n \sum_{i=1}^r (M_{ij} - M_j)^2}{n(r-1)}} = \sqrt{\frac{3.80029}{12 \times (7-1)}} = 0.2297$$

$$\text{CWS-Index for Expert - 4} = \frac{0.3742}{0.2297} = 1.629$$

The CWS-index for all experts are represented in Table-2. CWS-Index for Expert-1, Expert-2, Expert-3, Expert-4 and Expert-5 subsequently is 3.580, 8.237, 12.610, 1.629 and 7.612. Based on these CWS-Indexes, the Expertise-based Experts ranking obtained is [47]: Expert 3 - 2 - 5 - 1 - 4.

**Table-1.** The CWS-Index calculation for Expert-4.

Element Matrix	Original data	Formula-1		Formula-2		Formula-3		$M_j$	$r(M_j - GM)^2$	$\sum_{i=1}^r (M_{ij} - M_j)^2$
$p_{12}$	0.43	0.714	0.857	0.714	0.857	0.714	0.857	0.724	0.32655	0.13703
$p_{13}$	0.64	0.357	1	0.357	1	0.357	1	0.633	0.10789	0.72886
$p_{14}$	0.71	0.286	0.357	0.286	0.357	0.286	0.357	0.357	0.16037	0.16327
$p_{21}$	0.57	0.286	0.143	0.286	0.143	0.286	0.143	0.296	0.31635	0.16764
$p_{23}$	0.43	0.714	0.643	0.714	0.643	0.714	0.643	0.643	0.12636	0.11224
$p_{24}$	0.36	0.786	0.143	0.786	0.143	0.786	0.143	0.439	0.03403	0.77988
$p_{31}$	0.36	0.643	0	0.643	0	0.643	0	0.388	0.10206	0.68222
$p_{32}$	0.57	0.286	0.357	0.286	0.357	0.286	0.357	0.367	0.13948	0.05539
$p_{34}$	0.21	0.571	0.429	0.571	0.429	0.571	0.429	0.480	0.00585	0.11953
$p_{41}$	0.29	0.714	0.643	0.714	0.643	0.714	0.643	0.673	0.19050	0.13120
$p_{42}$	0.64	0.214	0.143	0.214	0.143	0.214	0.143	0.571	0.02772	0.63265
$p_{43}$	0.79	0.429	0.571	0.429	0.571	0.429	0.571	0.531	0.00342	0.09038
Total									1.54057	3.80029

**Table-2.** Discrimination, Inconsistency, CWS-Index and Experts Ranking.

	$e_1$	$e_2$	$e_3$	$e_4$	$e_5$
Discrimination	0.3703	0.3302	0.5835	0.3742	0.4662
Inconsistency	0.1035	0.0401	0.0463	0.2297	0.0612
CWS-Index	3.580	8.237	12.610	1.629	7.612
Rank	4	2	1	5	3

**Table-3.** The Importance Weights Calculation Using BUM  $Q(R) = R^\alpha$ .

	$e_3$	$e_2$	$e_5$	$e_1$	$e_4$
CWS- Index	12,6100	8,2370	7,6120	3,5800	1,6290
Log(CWS-Index)	1,1007	0,9158	0,8815	0,5539	0,2119
Accumulated(Log(CWS-Index))	1,1007	2,0165	2,8980	3,4519	3,6638
$R = \text{Normalized}(\text{Accumulated}(\text{Log}(\text{CWS-Index})))$	0,3004	0,5504	0,7910	0,9422	1,0000
$Q(R) = R^\alpha = R^{0.25}$	0,7403	0,8613	0,9431	0,9852	1,0000
Importance Weights for $\alpha = 0.25$ (in %)	74,03	12,10	8,17	4,22	1,48
$Q(R) = R^\alpha = R^{0.5}$	0,5481	0,7419	0,8894	0,9706	1,0000
Importance Weights for $\alpha = 0.5$ (in %)	54,81	19,38	14,75	8,13	2,94
$Q(R) = R^\alpha = R$	0,3004	0,5504	0,7910	0,9422	1,0000
Importance Weights for $\alpha = 1$ (in %)	30,04	25,00	24,06	15,12	5,78
$Q(R) = R^\alpha = R^2$	0,0903	0,3029	0,6256	0,8877	1,0000
Importance Weights for $\alpha = 2$ (in %)	9,03	21,27	32,27	26,20	11,23

Table-3 shows the importance weights calculation using BUM  $Q(R) = R^\alpha$ . After obtaining the rank, the experts are sorted by their expertise level. Expert-3 ( $e_3$ ) is an expert with the highest rank and placed on the first order. Expert-2 ( $e_2$ ) is in the second order. Then we calculate the logarithm of the CWS-Index and the

accumulation of the logarithm of the CWS-Index. After normalized the accumulation of the logarithm of the CWS-Index, the maximum value of these logarithm is 1 in accordance with the BUM function which has a maximum value of 1 and the experts total Importance weights are 1. In Table-3, the BUM function will be determined by





several values of the parameter  $\alpha$ . For example, for the parameters  $\alpha = 0.25$ ,  $\alpha = 0.5$ ,  $\alpha = 1$ ,  $\alpha = 2$ .

Suppose, we want to determine the importance weight of DMs for parameter  $\alpha = 0.25$ .

The CWS-Index for  $e_3$  is 12,610

For  $e_3$ ,  $\text{Log}(\text{CWS-Index}) = \log(12,610) = 1,1007$

For  $e_2$ ,  $\text{Log}(\text{CWS-Index}) = \log(8,2370) = 0,9158$

For  $e_3$ ,  $\text{Acc}((\text{Log}(\text{CWS-Index}))) = 1,1007$

For  $e_2$ ,  $\text{Acc}((\text{Log}(\text{CWS-Index}))) = 1,1007 + 0,9158 = 2,0165$

For  $e_3$ ,  $R = \text{Norm}(\text{Acc}(\text{Log}(\text{CWS-Index}))) = 1,1007/3,6638 = 0,3004$

For  $e_2$ ,  $R = \text{Acc}((\text{Log}(\text{CWS-index}))) = 2,0165/3,6638 = 0,5504$

If parameter  $\alpha = 0,25$ ,  $Q(R) = R^\alpha = R^{0.25}$

For  $e_3$ ,  $Q(0,3004) = 0,3004^{0.25} = 0,7405$

For  $e_2$ ,  $Q(0,3004) = 0,5504^{0.25} = 0,8613$

The Importance weight for  $e_3$  is 74, 05 %.

The Importance weight for  $e_2$  is 86, 13% - 74, 05 % = 12, 10 %

If parameter  $\alpha = 2$ , mismatches occur since the expert with higher expertise level gained smaller importance weight, for example Expert-3 ( $e_3$ ) as the expert with the highest expertise level, gets the smallest importance weight. If the parameter  $\alpha = 1$ , expert with higher level of expertise gained greater importance weight but not significantly. If the parameter  $\alpha = 0.5$ , expert with higher level of expertise gained greater importance weight significantly. Thus the parameter  $\alpha$  should be less than 1 to obtain the expected result, the higher the expert's expertise level, the higher his/her importance weight.

## CONCLUSIONS

An *expertise-based experts importance weight* method is proposed in order to develop the experts importance weight in adverse judgment situation in which every expert provides his/her judgment in pairwise comparisons FPR.

This model consists of 2 stages. The first stage, we obtain the experts' ranking by combining the experts' expertise level and Additive Consistency of FPR. In the second stage, we develop the experts' importance weight by using Basic Unit-Interval Increasing Monotonic Functions  $Q(R) = R^\alpha$ ,  $\alpha < 1$  to get the expected results, the higher the expert's expertise level, the higher his/her importance weight.

## ACKNOWLEDGEMENTS

Preparation of this paper was supported, in part, by the financing of Research Grants for Private University Lecturer in Kopertis Region 7, Fiscal Year 2014, No. 004/SP2H/P/K7/KM/2014, April 3, 2014 issued by Ministry of Education and Culture, Directorate General of Higher Education and the Research Funding issued by University of Surabaya.

## REFERENCES

- [1] V. Malhotra, M. D. Lee and A. Khurana. 2007. Domain experts influence decision quality: Towards a robust method for their identification. *Journal of Petroleum Science and Engineering*. 57(1-2): 181-194.
- [2] M. T. H. Chi, R. Glaser and M. J. Farr. 1988. *The Nature of Expertise*. New Jersey. Lawrence Erlbaum Associates.
- [3] J. R. Anderson. 2000. *Cognitive Psychology and its Implications*. Vol. 5. New York: Worth Publishing.
- [4] C. Horn. 2006. How experience affects perception in Expert Decision-Making. *Perception*. 05-814(2): 1-10.
- [5] R. M. Cooke. 1991. *Experts in uncertainty. Opinion and subjective probability in science*. New York/Oxford: Oxford University Press.
- [6] J. Shanteau, D. J. Weiss, R. P. Thomas and J. C. Pounds. 2002. Performance-based assessment of expertise: How to decide if someone is an expert or not? *European Journal of Operational Research*. 136(2): 253-263.
- [7] D. J. Weiss and J. Shanteau. 2003. Empirical Assessment of Expertise. *Human Factors*. 45(1): 104-116.
- [8] C. Yan, L. Jing, C. Xia and L. Yuanyuan. 2009. Method on the Assessment Level of Experts in Group Decision Making Based on Multi-granularity Linguistic Judgment Matrices. In *Information Engineering and Computer Science*. Wuhan. pp. 1-5.
- [9] E. Herowati, U. Ciptomulyono, J. Parung and Suparno. 2013. Competent-based Experts' Ranking at Fuzzy Preference Relations on Alternatives. In *International Conference on Industrial Engineering and Service Science*, Surabaya, Indonesia, pp. K7.1-K7.6.
- [10] R. L. Keeney. 1976. A group preference axiomatization with cardinal utility. *Management Science*. 23(2): 140-145.
- [11] F. Herrera, E. Herrera-Viedma and J. L. Verdegay. 1998. Choice Processes for Non-Homogeneous Group Decision Making in Linguistic Setting. *Fuzzy Sets and System*. 94(3): 287-308.
- [12] E. Charmela, J. Parung and E. Herowati. 2009. Pengembangan Model Integrasi Informasi Subjektif dan Objektif pada Multiple Attribute Group Decision Making. *Journal of Logistics and Supply Chain Management*. 2(2): 68-81.



- [13] S. E. Bodily. 1979. A Delegation Process for Combining Individual Utility Functions. *Management Science*. 25(10): 1035-1041.
- [14] R. Ramanathan and L. S. Ganesh. 1994. Group preference aggregation methods employed in AHP: An evaluation and an intrinsic process for deriving members' weightages. *European Journal of Operational Research*. 79(2): 249-265.
- [15] R. C. V. D. Honert. 2001. Decisional Power in Group Decision Making: A Note on the Allocation of Group Members' Weights in the Multiplicative AHP and SMART. *Group Decision and Negotiation*. 10: 275-286.
- [16] J. Kacprzyk, M. Fedrizzi and H. Nurmi. 1992. Group decision making and consensus under fuzzy preferences and fuzzy majority. *Fuzzy Sets System*. 49: 21-31.
- [17] D. Ben-Arieh and Z. Chen. 2006. Linguistic-Labels Aggregation and Consensus Measure for Autocratic Decision Making Using Group Recommendations. *IEEE Transactions on systems, man and cybernetics*. 36(3): 558-568.
- [18] R. O. Parreiras, P. Y. Ekel, J. S. C. Martini and R. M. Palhares. 2010. A Flexible Consensus Scheme for Multicriteria Group Decision Making Under Linguistic Assessments. *Information Sciences*. 180(7): 1075-1089.
- [19] Z. Xu. 2010. Interactive group decision making procedure based on uncertain multiplicative linguistic preference relations. *Journal of Systems Engineering and Electronics*. 21(3): 408-415.
- [20] Z. Xu and X. Cai. 2012. Minimizing Group Discordance Optimization Model for Deriving Expert Weights. *Group Decision and Negotiation*. 21(6): 863-875.
- [21] Z. Xu. 2008. Group Decision Making Based on Multiple Types of Linguistic Preference Relations. *Information Sciences*. 178(2): 452-467.
- [22] Z. L. Yue. 2011. An extended TOPSIS for determining weights of decision makers with interval numbers. *Knowledge-Based Systems*. 24(1): 146-153.
- [23] Z. L. Yue. 2012. Approach to group decision making based on determining the weights of experts by using projection method. *Applied Mathematical Modelling*. 36(7): 2900-2910.
- [24] Z. L. Yue. 2012. Extension of TOPSIS to determine weight of decision maker for group decision making problems with uncertain information. *Expert Systems with Applications*. 39(7): 6343-6350.
- [25] Z. L. Yue. 2011. A method for group decision-making based on determining weights of decision makers using TOPSIS. *Applied Mathematical Modelling*. 35(4): 1926-1936.
- [26] Z. L. Yue. 2011. Deriving decision maker's weights based on distance measure for interval-valued intuitionistic fuzzy group decision making. *Expert Systems with Applications*. 38(9): 11665-11670.
- [27] Z. L. Yue. 2012. Developing a straightforward approach for group decision making based on determining weights of decision makers. *Applied Mathematical Modelling*. 36: 4106-4117.
- [28] E. Herrera-Viedma, F. Chiclana, F. Herrera and S. Alonso. 2007. Group Decision-Making Model with Incomplete Fuzzy Preference Relations Based on Additive Consistency. *IEEE Transactions on Systems, Man, and Cybernetics (B)*. 37(1): 176-189.
- [29] S. Alonso, F. J. Cabrerizo, F. Chiclana, F. Herrera and E. Herrera-Viedma. 2009. Group Decision Making With Incomplete Fuzzy Linguistic Preference Relations. *International Journal of Intelligent Systems*. 24(2): 201-222.
- [30] X. Zhang, G. Yue and X. Zheng. 2011. Assigning Method for Decision Power Based on Linguistic 2-tuple Judgment Matrices. *Journal of Software*. 6(3): 508-515.
- [31] N. R. Ortiz, T. A. Wheeler, R. J. Breeding, S. Hora, M. A. Meyer and R. L. Keeney. 1991. Use of Expert Judgment in NUREG-1150. *Nuclear Engineering and Design*. 126(3): 313-331.
- [32] J. Kacprzyk. 1986. Group decision making with a fuzzy linguistic majority. *Fuzzy Sets and Systems*. 18(2): 105-118.
- [33] F. Chiclana, F. Herrera and E. Herrera-Viedma. 1998. Integrating three representation models in fuzzy multipurpose decision making based on fuzzy preference relations. *Fuzzy Sets and Systems*. 97(1): 33-48.
- [34] E. Herrera-Viedma, F. Herrera, F. Chiclana and M. Luque. 2004. Some Issues on Consistency of Fuzzy Preference Relations. *European Journal of Operational Research*. 154(1): 98-109.
- [35] S. A. Orlovski. 1978. Decision-making with fuzzy preference relations. *Fuzzy Sets and Systems*. 1(3): 155-167.



- [36] T. Tanino. 1984. Fuzzy Preference Orderings in Group Decision Making. *Fuzzy Sets and Systems*. 12(2): 117-131.
- [37] F. Chiclana, E. Herrera-Viedma and S. Alonso. 2009. A note on two methods for estimating missing pairwise preference values. *IEEE Transactions on systems, man and cybernetics Part B*. 39(6): 1628-1633.
- [38] E. Herrera-Viedma, S. Alonso, F. Chiclana and F. Herrera. 2007. A Consensus Model for Group Decision Making With Incomplete Fuzzy Preference Relations. *IEEE Transactions on Fuzzy Systems*. 15(5): 863-877.
- [39] S. Alonso, F. Chiclana, F. Herrera, E. Herrera-Viedma, J. A. 'a-Fdez and C. Porcel. 2008. A Consistency-Based Procedure to Estimate Missing Pairwise Preference Values. *International Journal of Intelligent Systems*. 23(2):155-175.
- [40] G. Zhang, Y. Dong and Y. Xu. 2012. Linear optimization modeling of consistency issues in group decision making based on fuzzy preference relations. *Expert Systems with Applications*. 39(3): 2415-2420.
- [41] S. Alonso, E. Herrera-Viedma, F. Chiclana and F. Herrera. 2010. A web based consensus support system for group decision making problems and incomplete preferences. *Information Sciences*. 180(23): 4477-4495.
- [42] F. Chiclana, E. Herrera-Viedma, F. Herrera and S. Alonso. 2007. Some induced ordered weighted averaging operators and their use for solving group decision making problems based on fuzzy preference relations. *European Journal of Operational Research*. 182(1): 383-399.
- [43] R. R. Yager. 1988. On ordered weighted averaging aggregation operators in multi-criteria decision making. *IEEE Transactions on Systems, Man, and Cybernetics*. 18(1): 183-190.
- [44] R. R. Yager and D. P. Filev. 1999. Induced ordered weighted averaging operators. *IEEE Transaction Systems, Man, and Cybernetics, Part B: Cybernetics*. 29(2): 141-150.
- [45] R. R. Yager. 2004. OWA Aggregation over a Continuous Interval Argument with Applications to Decision Making. *IEEE Transactions on Systems, Man and Cybernetics part B: Cybernetics*. 34(5): 1952-1963.
- [46] R. R. Yager. 1996. Quantifier Guided Aggregation Using OWA Operators. *International Journal of Intelligent Systems*. 11: 49-73, 1996.
- [47] E. Herowati, U. Ciptomulyono, J. Parung and Suparno. Expertise-based Experts Ranking in Adverse Judgment. Working in process paper, vol. -.